

学校编码: 10384

分类号_____密级_____

学号: X200326008

UDC _____

厦 门 大 学

硕 士 学 位 论 文

香港海域浮游植物种类多样性与群落动态
研究

Species Diversity and Community Dynamics of
Phytoplankton in Coastal Waters of Hong Kong

林 旭 吟

指导教师姓名: 高亚辉 教授

专 业 名 称: 微生物学

论文提交日期: 2007 年 9 月

论文答辩时间: 2007 年 9 月 22 日

学位授予日期:

答辩委员会主席: 何建宗教授

评 阅 人: _____

2007 年 9 月

厦门大学博硕士论文摘要库

厦门大学学位论文原创性声明

兹呈交的学位论文，是本人在导师指导下独立完成的研究成果。本人在论文写作中参考的其他个人或集体的研究成果，均在文中以明确方式标明。本人依法享有和承担由此论文产生的权利和责任。

声明人（签名）：

年 月 日

厦门大学博硕士论文摘要库

厦门大学学位论文著作权使用声明

本人完全了解厦门大学有关保留、使用学位论文的规定。厦门大学有权保留并向国家主管部门或其指定机构送交论文的纸质版和电子版，有权将学位论文用于非赢利目的的少量复制并允许论文进入学校图书馆被查阅，有权将学位论文的内容编入有关数据库进行检索，有权将学位论文的标题和摘要汇编出版。保密的学位论文在解密后适用本规定。

本学位论文属于

1、保密（ ），在 年解密后适用本授权书。

2、不保密（ ）

（请在以上相应括号内打“√”）

作者签名：

日期： 年 月 日

导师签名：

日期： 年 月 日

厦门大学博硕士论文摘要库

目录

中文摘要	I
Abstract.....	I
第一章 绪论	1
1. 1 香港地理概况.....	1
1. 2 香港海域环境介绍.....	2
1. 3 香港海域污染情况.....	3
1. 4 香港海域的赤潮.....	5
1. 5 香港海域浮游植物研究动态.....	7
1. 6 香港海域域浮游植物监测计划.....	8
1. 7 本论文研究目标和内容.....	9
第二章 材料与方法	11
2. 1 采样站位介绍.....	11
2. 2 样品的采集、处理和观察.....	12
2. 3 主要仪器设备.....	13
2. 4 数据处理及统计分析.....	13
2. 4. 1 群落结构分析.....	13
2. 4. 2 数据统计分析.....	14
2. 5 藻类分类鉴定所用书籍.....	14
第三章 结果与分析	15
3. 1 香港海域不同季节理化环境因子的变化.....	15
3. 1. 1 香港海域表层海水温度变化.....	15
3. 1. 2 香港海域表层海水盐度变化.....	15
3. 1. 3 香港海域表层海水溶解氧状况.....	16
3. 1. 4 香港海域表层海水 5 天生化耗氧量（5-BOD）的状况.....	17
3. 1. 5 香港海域表层海水营养盐状况.....	19
3. 2 香港海域浮游植物群落结构特征与动态.....	24
3. 2. 1 香港海域浮游植物种类组成.....	24

3. 2. 2 香港海域浮游植物优势种组成	25
3. 2. 3 香港海域浮游植物优势属、种的季节变化	27
3. 2. 4 香港海域浮游植物数量的季节变化	30
3. 2. 5 香港海域浮游植物群落组成的季节变化	30
3. 2. 6 香港海域浮游植物种类数的季节变化	33
3. 2. 7 香港海域不同站位浮游植物多样性指数的分布	33
3. 2. 8 香港海域不同站位浮游植物细胞密度的分布	34
3. 2. 9 香港海域浮游植物数量、叶绿素 a 的含量与环境因子关系	37
第四章 讨论	38
4. 1 2000~2004 年香港海域的温、盐、营养盐特征	38
4. 2 2000~2004 年香港海域浮游植物群落特征	39
4. 2. 1 香港海域浮游植物的空间分布特征	39
4. 2. 2 香港海域浮游植物优势种变化特征	39
4. 2. 3 香港不同海域浮游植物组成特征	40
4. 2. 4 香港海域浮游植物群落结构的评价	43
4. 3 2000~2004 年香港海域浮游植物群落动态与环境因子的关系	43
4. 3. 1 盐度对香港海域浮游植物的影响	43
4. 3. 2 硅对香港海域浮游植物的影响	44
4. 3. 3 TN/TP 值对香港海域浮游植物的影响	44
第五章 结论与展望	46
5. 1 结论	46
5. 1. 1 香港海域理化环境因子的变化	46
5. 1. 2 香港海域浮游植物群落的特征	46
5. 2 本论文的特色	47
5. 3 本论文的不足之处和今后的工作展望	47
5. 3. 1 本论文的不足之处	47
5. 3. 2 今后的工作展望	48
参考文献	49
附表	57

图版	82
攻读硕士阶段发表的论文和参加的课题	94
致谢	95

厦门大学博硕士论文摘要库

Contents

Abstract.....	I
Chapter I Introduction.....	1
1. 1 Geography survey of HongKong Island.....	1
1. 2 Environmental character of HongKong coastal areas.....	2
1. 3 Pollution of Hongkong coastal areas	3
1. 4 Red tides in Hongkong coastal areas	5
1. 5 Study of phytoplankton in Hong Kong waters	7
1. 6 Supervision about phytoplankton in Hongkong coastal areas	8
1. 7 Purpose and main content of the present study	9
Chapter II Materials and methods.....	11
2. 1 Sampling stations	11
2. 2 Sampling, processing and observation of samples	12
2. 3 Instruments and equipments	13
2. 4 Data treatment and statistical analyses	13
2. 4. 1 Community structure analyses.....	13
2. 4. 2 Statistical analyses.....	14
2. 5 References for taxonomy	14
Chapter III Results and Analysis	15
3. 1 Environmental factors in different seasons in coastal waters of Hongkong	15
3. 1. 1 Temperature characteristics of surface sea water.....	15
3. 1. 2 Salinity characteristics of surface sea water	15
3. 1. 3 Dissolved oxygen characteristics of surface sea water.....	16
3. 1. 4 5-BOD in surface sea water.....	17
3. 1. 5 Nutrients characteristics of surface sea water.....	19
3. 2 Characteristics of phytoplankton community structure in coastal waters of Hongkong.....	24

3. 2. 1	Phytoplankton composition.....	24
3. 2. 2	Dominant species composition.....	25
3. 2. 3	Seasonal change of dominant phytoplankton genus and species..	27
3. 2. 4	Seasonal change of phytoplankton species numbers.....	30
3. 2. 5	Seasonal change of phytoplankton cell density.....	30
3. 2. 6	Seasonal change of phytoplankton species composition.....	33
3. 2. 7	Diversity indexes of phytoplankton species at different stations .	33
3. 2. 8	Distribution of phytoplankton abundance at different stations	34
3. 2. 9	Relationship between phytoplankton abundance, Chl <i>a</i> and environmental factors.....	37
Chapter IV Conclusion.....		38
4. 1	Temperature, salinity and nutrients charactersitics in 2000-2004	38
4. 2	Charactersitics of phytoplankton community in 2000-2004.....	39
4. 2. 1	Temporal and spatial distribution of phytoplankton.....	39
4. 2. 2	Variation of dominant diatom species	39
4. 2. 3	Composition of phytoplankton species in different sea areas	40
4. 2. 4	Evaluation on ecological character of phytoplankton community	43
4. 3	Relationship between phytoplankton and environmental factors in 2000-2004	43
4. 3. 1	Influence of salinity on phytoplankton	44
4. 3. 2	Influence of silicate on phytoplankton.....	44
4. 3. 3	Influence of TN/TP on phytoplankton	46
Chapter V Conclusion and Prospect.....		46
5. 1	Conclusion	46
5. 1. 1	Environmental fators.....	46
5. 1. 2	Character of phytoplankton community	46
5. 2	Main innovatives of the dissertation	47
5. 3	Problems to be further studied	47

5. 3. 1 Problems of the dissertation	47
5. 3. 2 Further studies	48
Reference	49
Appidex table	57
Figureplate	82
Acknowledgements	95

中文摘要

香港海域是我国的赤潮高发区之一，浮游植物的长期监测是海洋环境调查、海水水质监测、赤潮研究的基础。本论文通过对 2000 年、2002 年和 2004 年，分春、夏、秋、冬四个季节在香港海域采集的浮游植物样品的种类鉴定分析，对香港海域浮游植物的种类多样性、种群分布特征和群落组成结构与动态及其与环境因子的关系等进行了研究，并从生态学角度探讨了香港海域浮游植物群落结构及变动的规律和主要机制。

结果如下：

1. 香港海域的盐度在平面分布上有显著的地理差异。根据受珠江口淡水的影响程度，将香港海域分成东部、中部、南部和西部水域，受珠江口淡水影响的程度由西向东逐渐减弱，形成了西部河口性质的水域和东部海洋性质的水域，使不同水域出现了不同的浮游植物群落结构特征。

2. 香港海域营养盐水平整体较低，总无机氮和总氮变化范围分别为 0.01~0.96mg/L、0.06~1.17 mg/L，正磷酸盐和总磷变化范围分别为 0.002~0.079mg/L、0.02~0.16 mg/L，硅含量的变化范围是 0.05~3.3 mg/L。与 20 世纪 80 年代末的数据比较有显著下降，而与 1999 年的数据比较则相对稳定。

3. 本研究共鉴定到浮游植物 254 种（含变种、变型），其中硅藻 51 属 177 种，甲藻 22 属 64 种，硅鞭藻 3 属 4 种，裸藻 1 属 1 种，针胞藻 2 属 2 种，金藻 1 属 1 种，隐藻 1 属 2 种，蓝藻 1 属 2 种，绿藻 1 属 1 种。硅藻仍然是香港海域浮游植物的主要类群，占总种数的 69.69%。其中，出现种类最多的是角毛藻属（*Chaetoceros*），有 27 种，占 10.63%；但在细胞密度上以骨条藻为优势，占了总细胞密度的 44.69%。

4. 与历史资料比较，香港海域浮游植物优势种有一定的变化，由原来的硅藻绝对优势转变为在一定季节和区域出现了甲藻或隐藻占优势的情况。2000 年春季的优势种是伸长斜片藻（*Plagioselmis prolunga*），而夏、秋、冬三季节以硅藻为优势，主要优势种有骨条藻（*Skeletonema* spp.）、微小细柱藻（*Leptocylindrus minimus*）、冰河星杆藻（*Asterionellopsis glacialis*）。2002 年的春季和夏季主要的优势种是骨条藻（*Skeletonema* spp.），而在秋季则出现以微小原甲藻（*Prorocentrum minimum*）为主要优势种，冬季的优势种是冰河星杆藻（*Asterionellopsis glacialis*）

和新月筒柱藻 (*Cylindrotheca closterium*)。2004 年的春季、夏季和秋季仍然以硅藻为优势, 主要的优势种有旋链角毛藻 (*Chaetoceros curvisetus*)、骨条藻 (*Skeletonema* spp.)、拟菱形藻 (*Pseudo-nitzschia* spp.) 和脆根管藻 (*Rhizosolenia fragillissima*), 但冬季的优势为微小原甲藻 (*Prorocentrum minimum*)。

5. 香港海域浮游植物数量的季节变化显著, 一般春季和冬季的细胞密度较低, 细胞密度的高峰均出现在夏季和夏末秋初。2000 年、2002 年和 2004 年的年平均细胞密度分别为 $2.65 \times 10^5 \text{ cells/L}$ 、 $6.90 \times 10^5 \text{ cells/L}$ 和 $7.73 \times 10^5 \text{ cells/L}$, 呈现出逐年增加的趋势。香港海域浮游植物细胞密度因受珠江口淡水的影响程度不同, 在平面分布上表现出一定的差异。东部水域的浮游植物细胞密度季节变化不明显, 而中部、南部和西部水域则有明显的季节变化, 通常在夏季大量径流注入时形成细胞密度的高峰。

6. 从群落组成来看, 东部水域甲藻和其他藻类所占比例相对高于其他三个水域; 而中部水域、南部水域和西部水域多数季节以硅藻为优势, 但其他藻类的比例高于甲藻。

7. 2000 年、2002 年和 2004 年香港海域浮游植物种类多样性指数分别为 2.56、2.54 和 2.17, 逐年略呈递减的趋势但总体较稳定。调查期间, 群落的多样性指数平均值为 2.42, 均匀度平均值为 0.54, 属于中等水平。

8. 对香港海域浮游植物细胞密度和叶绿素 a 含量与环境因子 (温度、盐度、亚硝酸盐、硝酸盐、氨氮、总氮、总磷、硅 Si、TN/TP、Si/TN、Si/TP) 进行了逐步回归, 结果表明, 香港海域浮游植物细胞密度和叶绿素 a 含量与盐度、硅含量和 TN/TP 值相关性极其显著, 说明盐度、硅含量和 TN/TP 值是影响香港海域浮游植物生物量的主要环境因素。

关键词: 香港海域; 浮游植物; 种类; 赤潮; 生态

Abstract

Hong Kong is one of the coastal areas with frequent red tide events, and long-term monitor for phytoplankton is the foundation for marine environment investigation, water quality control and the red tide studies. In the present studies, various phytoplankton samples were collected and analyzed from the coastal waters of Hong Kong, in four seasons of 2000, 2002 and 2004. Through the species identification of these phytoplankton samples, our studies have been focused on investigating phytoplankton species composition and diversity, population distribution, community structure and dynamics in the relationship with the environmental factors of Hong Kong coastal waters. Based on these studies, we proposed from a view of ecology the main mechanism of phytoplankton community structure as well as the dynamics in Hong Kong coastal waters. The results are as follow:

1. There are significant geographical differences in the horizontal distribution of salinity in Hong Kong coastal waters. Hong Kong coastal area can be divided into eastern, central, southern and western regions, based on different impact from the freshwater of Pearl River. The salinity is gradually weakened from west to east, leading to the formation of limnetic western region and oceanic eastern region. Therefore the different groups of phytoplankton changed in their structural characteristics according to the different water regions.
2. The overall level of nutrients in Hong Kong coastal waters is relative low. Total nitrogen and total inorganic nitrogen varied from 0.01-0.96 mg/L、0.06-1.17 mg/L, respectively. Phosphate and phosphorous varied from 0.002-0.079 mg/L、0.02-0.16 mg/L, respectively. The silicon content is in the range of 0.05-3.3 mg/L. The nutrient level was dramatically reduced compared to that of the end 1980s, but is relative stable compared to that of 1999.
3. Totally, 254 species (including varieties) of phytoplankton were identified. They are including 177 diatom species belonging to 51 genus, 64 dinoflagellate species belonging to 22 genus, 4 silicoflagellate species belonging to 3 genus, 1 englenoid species belonging to 1 genus, 2 raphidophyte species belonging to 2 genus, 1

chrysophyte species belonging to 1 genus, 1 cryptomonad species belonging to 2 genus, 1 Cyanophyta species belonging to 2 genus, 1 Chloophyta species belonging to 1 genus. In the waters of Hong Kong, diatoms is the major component of phytoplankton, accounting for of 69.69% of the total species numbers of phytoplankton. Among these, the most frequently appearance diatom genus is *Chaetoceros* (27 species, representing 10.63%); while according to the cell densities, *Skeletonema* is the largest, about 44.69% of total phytoplankton cell densities.

4. Comparing to literature records, the dominant phytoplankton species of Hong Kong coastal waters changed to some extent. Diatom had been reported as the most predominant group in Hong Kong coastal waters. In this study it was shown that in certain seasons, Dinoflagellate or Cryptomonad appeared to be dominant in some sea areas. In the spring of 2000, the dominant species was *Plagioselmis prolunga* (Cryptomonad), while in summer, autumn and winter, the diatom dominated Hong Kong coastal waters, with the dominant species of *Skeletonema* spp., *Leptocylindrus minimus*, and *Asterionellopsis glacialis*. In the spring and summer of 2002, the dominant species was *Skeletonema* spp., while in autumn it was *Prorocentrum minimum*, and in winter the dominant species were *Asterionellopsis glacialis* and *Cylindrotheca closterium*. In the spring, summer and autumn of 2004, diatom still dominated in Hong Kong coastal waters, with the dominant species of *Chaetoceros curvisetus*, *Skeletonema* spp., *Pseudo-nitzschia* spp. and *Rhizosolenia fragillissima*, while in winter Dinoflagellate was the dominant group and the dominant species was *Prorocentrum minimum*.

5. The abundance of phytoplankton showed significant season dependent variation. Generally cell density of phytoplankton appeared lower in spring and winter, but higher in summer and early autumn. In 2000, 2002 and 2004, the annual average of cell density of phytoplankton increased, with the average cell density of 2.65×10^5 cells/L, 6.90×10^5 cells/L, and 7.73×10^5 cells/L, respectively. The horizontal distribution of phytoplankton cell density varied in different sea areas according to the extent of effect exerted by fresh water from Pearl River. Cell density did not fluctuate obviously by seasons in eastern waters, while in central, southern and western waters,

Degree papers are in the "[Xiamen University Electronic Theses and Dissertations Database](#)". Full texts are available in the following ways:

1. If your library is a CALIS member libraries, please log on <http://etd.calis.edu.cn/> and submit requests online, or consult the interlibrary loan department in your library.
2. For users of non-CALIS member libraries, please mail to etd@xmu.edu.cn for delivery details.

厦门大学博硕士论文摘要库